

Homework 6

1. *A Hovercraft moving under random wind gusts*

Here we will consider a hovercraft with mass 1 moving under a random wind gusts. We will use an extremely simplified model of wind gusts. We'll see how to improve this model later in the course.

We have a hovercraft moving in the plane with two thrusters, each pointing through the center of mass, exerting forces in the \mathbf{x} and \mathbf{y} directions. The hovercraft has mass 1. The discretized equations of motion for the hovercraft are

$$x(t+1) = \begin{bmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{bmatrix} x(t) + \begin{bmatrix} \frac{1}{2} & 0 \\ 1 & 0 \\ 0 & \frac{1}{2} \\ 0 & 1 \end{bmatrix} \begin{bmatrix} u_1(t) \\ u_2(t) \end{bmatrix}$$

where x_1 and x_2 are the position and velocity in the \mathbf{x} -direction, and x_3, x_4 are the position and velocity in the \mathbf{y} -direction. Here

$$u(t) = \begin{bmatrix} u_1(t) \\ u_2(t) \end{bmatrix}$$

is the force acting on the hovercraft for time in the interval $[t, t+1)$.

Suppose the forces acting on the hovercraft are i.i.d. and Gaussian, with $u(t) \in \mathcal{N}(0, \Sigma_u)$, and $u(s)$ and $u(t)$ uncorrelated when $s \neq t$, and

$$\Sigma_u = \begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix}$$

- Find an expression for the 90% confidence ellipsoid for the position of the hovercraft after 10 seconds. Plot the ellipsoid in \mathbb{R}^2 .
- Simulate the system 1000 times and plot the resulting final positions in the plane. Count (using Matlab) the percentage of final positions that land in the 90% confidence ellipsoid.
- Also plot the 90%, 70% and 50% confidence ellipsoids on the same plot.
- The final position $x \in \mathbb{R}^2$ is Gaussian, with mean zero. Suppose its covariance is Σ . What is the pdf of $x^T \Sigma^{-1} x$? Compute $x^T \Sigma^{-1} x$ for 1000 simulations and plot the empirical cdf on top of the actual cdf.

2. *The Chebyshev inequality for vectors and confidence ellipsoids*

- Suppose $x : \Omega \rightarrow \mathbb{R}$. Show that the Chebyshev inequality can be written as


$$\text{Prob}\left(\frac{(x - \mu)^2}{\text{cov}(x)} \geq a\right) \leq \frac{1}{a}$$

where $\mu = \mathbb{E}x$.

- Suppose $x : \Omega \rightarrow \mathbb{R}$ is a scalar random variable, which is uniformly distributed on the interval $[-1, 1]$. The Chebyshev bound gives a lower bound that x lies in the interval $[-c, c]$. What are the values of c corresponding to lower bounds of 0.1, 0.5 and 0.9?
- Now let's consider a random vector $x : \Omega \rightarrow \mathbb{R}^n$, with mean $\mathbb{E}x = \mu$ and covariance matrix $\text{cov}(x) = \Sigma$. Generalize the Chebyshev inequality to find an upper bound on the probability

$$\text{Prob}\left((x - \mu)^T \Sigma^{-1} (x - \mu) \geq a\right)$$

- (d) This can be used to construct *confidence ellipsoids* for pdfs which are not Gaussian. For example, consider the induced pdf for the random variable $x : \Omega \rightarrow \mathbb{R}^2$ which is uniform on the triangle shown below.



What is the mean and covariance of x ?

- (e) Using the Chebyshev bound you derived above, plot ellipsoids E_α such that

$$\text{Prob}(x \in E_\alpha) \geq \alpha$$

for the cases where α is 0.1, 0.5 and 0.9.

3. Conditional pdfs for Gaussians

Suppose $x \in \mathbb{R}^n$ and $x \sim \mathcal{N}(0, \Sigma)$. Partition x as $x = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$ where $x_1 \in \mathbb{R}^r$, and partition Σ as

$$\Sigma = \begin{bmatrix} \Sigma_{11} & \Sigma_{12} \\ \Sigma_{12}^T & \Sigma_{22} \end{bmatrix}$$

so that $\Sigma_{11} \in \mathbb{R}^{r \times r}$. First let's consider the special case when $n = 2$, $r = 1$ and

$$\Sigma = \begin{bmatrix} 2 & 0.8 \\ 0.8 & 1 \end{bmatrix}$$

- Plot the 90% confidence ellipsoid C for x .
- Suppose now we are given $x_2 = 1$. Find the 90% confidence interval $[c_1, c_2]$ for the conditional $x_1 | (x_2 = 1)$, and find its mean. Plot three vertical lines through the boundary points of this interval and the mean.
- Now collect 10000 samples of x . In Matlab, pick out those samples for which $0.9 \leq x_2 \leq 1.1$, and discard the rest. Count the percentage of those you kept for which x_1 lies in $[c_1, c_2]$.
- The ellipsoid C intersects the line $x_2 = 1$ at points x_1 in the interval $[c_3, c_4]$. Find this interval.
- More generally, suppose E is the ellipsoid

$$E = \left\{ x \in \mathbb{R}^n \mid x^T \Sigma^{-1} x \leq 1 \right\}$$

The intersection of E with the surface

$$\left\{ \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \in \mathbb{R}^n \mid x_2 = b \right\}$$

is an ellipsoid, which has the form

$$F = \left\{ x_1 \in \mathbb{R}^r \mid (x_1 - a)^T Q^{-1} (x_1 - a) \leq c \right\}$$

Find a , c and Q in terms of b and Σ .